

Exploratory Analysis of Spanish Energetic Mining Accidents

Lluís Sanmiquel

Dpto. de Ingeniería Minera y Recursos Naturales, Universitat Politècnica de Catalunya,
Barcelona, Spain

Modesto Freijo

Dpto. de Ingeniería Eléctrica, Universitat Politècnica de Catalunya, Barcelona, Spain

Josep M. Rossell

Dpto. de Matemática Aplicada III, Universitat Politècnica de Catalunya, Barcelona, Spain

Using data on work accidents and annual mining statistics, the paper studies work-related accidents in the Spanish energetic mining sector in 1999–2008. The following 3 parameters are considered: age, experience and size of the mine (in number of workers) where the accident took place. The main objective of this paper is to show the relationship between different accident indicators: risk index (as an expression of the incidence), average duration index for the age and size of the mine variables (as a measure of the seriousness of an accident), and the gravity index for the various sizes of mines (which measures the seriousness of an accident, too). The conclusions of this study could be useful to develop suitable prevention policies that would contribute towards a decrease in work-related accidents in the Spanish energetic mining industry.

energetic mining risk index average duration index gravity index mining accidents

1. INTRODUCTION

In the total number of Spanish economic sectors, mining is one of those with a high index of annual incidences, i.e., the number of accidents per 100000 workers. In 2006, Spanish mining had an incidence index 4.7 times higher than all economic sectors [1, 2, 3]. If the indices of work accidents in Spanish mining are compared with those of other countries, we can see that their values are also much higher. Specifically, in 2006, the index of annual incidence was 8.9 times higher than

in the USA [4] and 20.4 times higher than in the State of Queensland (Australia) [5]. The Spanish mining sector can be classified into two types, energetic and nonenergetic mining. Energetic mining includes activities related to extraction and agglomeration of coal, extraction of uranium and thorium, and extraction and preparation of any solid fuel. Petrol and gas extraction are not included. Nonenergetic mining includes the extraction and preparation of metallic and nonmetallic minerals and quarry products such as limestone, marble, granite, sand and gravel or clay, with the

We would like to thank the National Institute of Safety and Hygiene in Work for allowing us to use the annual general databases for accidents in Spain in the mining sector from 1999–2007. We also wish to thank the Director General of Energetic Policies and Mining of the Spanish Ministry of Industry, Tourism and Commerce for giving access to the mining statistics data from 1999–2006.

Correspondence and requests for offprints should be sent to Lluís Sanmiquel, Polytechnical University of Catalunya, Avda. Bases de Manresa, 61–73, 08242 Manresa (Barcelona), Spain. E-mail: sanmi@emrn.upc.edu.

purpose of producing aggregates, cement, ornamental stones, concrete or ceramic products. The indices of annual incidences in energetic mining [1] are significantly larger than in nonenergetic mining [2]. In 2006, this index was 2.7 times higher than in the nonenergetic sector.

Most accidents are caused by human error [6], and many of those are experienced by relatively few workers [7]. The first event that immediately precedes an accident is mostly caused by environment factors, while the second event, i.e., the event that takes place immediately before the first one, is mainly attributed to behaviour factors [8].

This paper considers several characteristics. Specifically, it analyses the relationship between work-related accidents and the age of workers, occupational experience and the size of the mining work centre (defined by the number of employees).

Many researchers studied the effect of age and mining experience on the occurrence of accidents; different conclusions followed. The National Research Council found a strong negative link between age and the seriousness of the lesions of injured workers in coal mines [9]. Younger injured workers seemed more seriously injured than the older ones. Bennett did not find any relationship between age and the seriousness of the injury; the author concluded that less experienced miners were slightly more likely to have an accident than more experienced ones [10]. However, Bennett and Passmore showed that older injured miners were more likely to have serious or fatal injuries than younger ones [11].

Butani noted that injuries in the coal industry were more related to experience than to age, and less experience increased the probability of an accident [12]. Specifically, this study concludes that there exists a significant occupational experience effect with the largest increase in risk occurring in workers over 50 years old with under one year of occupational experience. In their study on transport injuries in small coal mines, Hunting and Weeks reported an increased risk of injury with less experience but no age effect [13]. They also observed that small coal mines had the highest rates of transport-related injuries. Moreover, small mines had a greater share of fatal and

permanently disabling injuries, whereas the large mines had a greater share of injuries involving no lost workdays. On the other hand, some studies showed how accidents in the workplace could be attributed to personal and environmental factors [14, 15, 16, 17].

Groves, Kecojevic and Komljenovic classified accidents taking into account the machine or equipment used when the accident happened [18]. A similar study has been impossible with our available mining data. In this paper, information on the machine or equipment has been substituted with seven frequent types of accidents occurring in underground and surface mining.

We can note that a wide variety of results has been offered in the literature depending on the available mining data. In this article, we have presented and analysed data on accidents in the Spanish energetic mining sector in 1999–2008. The conclusions may be used in planning appropriate safety programs and measures such as engineering, enforcement, education or technological advances, to warn against injuries or fatal accidents in the energetic mining sector in the future.

2. METHODS

2.1. Study Population

The study population comprised accidents that took place in the Spanish energetic mining sector in 1999–2008, within the work schedule (we did not consider accidents which happened on one's way to or from work) which caused the injured worker to lose at least one workday.

The data were obtained from the annual digital database of the Ministry of Employment and Social Security with ArcGis 9.2¹. It is important that the annual accident database does not supply separate information about underground mining activities and surface ones for 1999–2002. However, this information is available for 2003–2008, and the two kinds of mining activities can be distinguished. In this paper, we analyse the overall accidents which happened in the indicated period or only the accidents corre-

sponding to 2003–2008, according to information from the database.

The percentage of workers in the Spanish energetic mining who were divided into seven age groups and into six types of work centres, were obtained from the yearbooks of work statistics of Spain's Ministry of Work and Immigration from 1999–2008 and the yearbooks of mining statistics of Spain's Ministry of Industry, Tourism and Commerce for 1999–2006.

2.2. Description of the Methodology

The study was divided into underground and surface mining, but this only applied to 2003–2008, since in 1999–2002 the location of the accidents was unknown (see section 2.1.).

We considered the workers' age and the size of the work centre (defined by the number of employees) where the accident took place. There were seven age groups (16–24, 25–29, 30–34, 35–39, 40–44, 45–54 and ≥ 55 years), six sizes of work centres (1–9, 10–19, 20–49, 50–99, 100–499 and ≥ 500 workers) and six experience groups (0–12, 13–60, 61–120, 121–180, 181–240 and ≥ 241 months) for seven kinds of accidents.

A risk index is defined as the ratio of the percentage of injuries attributed to a given subpopulation (age group or size of work centre) to the percentage of the total workforce represented by that subpopulation [12]. A risk index of 1 corresponds to an average risk, while a value greater than 1 indicates a higher risk for that group. Thus, to calculate the risk index, we needed to know the percentage of accidents that happened in each age groups or work centre, and the percentage of workers in them. We also calculated the average duration index (ADI) for each age group (1999–2007) and for each size of the work centre (1999–2006); it indicated the seriousness of accidents.

Once the risk index and the ADI for the age groups were calculated, we analysed the relationship between the risk index (as an indicator of the incidence of accidents in a population) and the ADI (as an indicator of the severity of the

accidents), and both the age and the size of the work centres. This was done with the nonparametric statistical Spearman rank correlation. The mean was calculated for the total population in 1999–2006 or 1999–2008. Throughout this paper, analyses were conducted at a .05 significance level.

The results were analysed in two ways. The Mann–Whitney *U* test was used to observe if there was a significant difference between underground and surface mining in the distribution of the ADI, depending on the occupational experience of the injured parties. Spearman rank correlation coefficient was used to analyse the relationship between the ADI and the workers' experience in the seven types of the most frequent accidents in underground and surface mining.

3. RESULTS AND DISCUSSION

3.1. Age and Experience of Injured Workers

Table 1 shows data on the percentage of workers in 1999–2008 per year, per age group, for the seven age groups. The results in Table 2 show a significantly greater risk in the 16–24, 25–29, 30–34 and 35–39 age groups, with the highest estimated risk indices. At the same time, the risk index was significantly lower for the 45–54 and, especially, ≥ 55 groups.

Spearman rank correlation coefficients, calculated for each year, show that the incidence of accidents (expressed with the risk index) is correlated with the workers' age: all coefficients are over the critical value of .714. The correlation is negative, i.e., when the workers' age increases, the incidence of accidents decreases (Table 2). Table 2 illustrates accidents for each age group, the risk index and Spearman rank correlation coefficient for each year between 1999 and 2008.

To analyse the possible relationship between the seriousness of the accidents and the age of the workers, the ADI for each age group was calculated with data from Table 3 (Table 4). Spearman rank correlation coefficients were higher than the critical value, except in 2005 and 2006. This indi-

¹ <http://www.esri.com/software/arcgis/arcgisserver/>

TABLE 1. Energetic Mining Population by Age Group (1999–2008)

Year	Age Group (%)						
	16–24	25–29	30–34	35–39	40–44	45–54	≥55
1999	3.09	7.73	19.07	31.96	26.29	9.79	2.06
2000	2.75	6.59	18.13	31.32	29.12	10.44	1.65
2001	2.41	6.02	16.87	30.72	30.72	10.84	2.41
2002	2.01	6.04	15.44	30.87	32.21	11.41	2.01
2003	2.24	5.97	14.18	30.60	32.84	12.69	1.49
2004	1.68	5.88	12.61	30.25	32.77	14.29	2.52
2005	1.92	5.77	11.54	29.81	32.69	15.38	2.88
2006	2.13	5.32	11.70	28.72	32.98	17.02	2.13
2007	2.33	5.81	11.63	26.74	32.56	18.60	2.33
2008	2.53	6.33	12.66	24.05	34.18	18.99	1.27
M	2.37	6.27	15.15	30.07	31.06	13.01	2.07

TABLE 2. Distribution of Accidents by Age Group and Estimated Risk Index (1999–2008)

Year	Age Group (%)							Risk Index							Spearman Correlation
	16–24	25–29	30–34	35–39	40–44	45–54	≥55	16–24	25–29	30–34	35–39	40–44	45–54	≥55	
1999	4.31	8.33	20.82	33.63	26.76	5.52	0.64	1.39	1.08	1.09	1.05	1.02	0.56	0.31	–.964
2000	4.41	8.04	20.59	31.67	29.09	5.54	0.66	1.60	1.22	1.14	1.01	1.00	0.53	0.40	–1.000
2001	2.58	6.74	19.03	34.04	32.15	4.87	0.59	1.07	1.12	1.13	1.11	1.05	0.45	0.25	–.750
2002	2.87	7.43	16.65	32.85	34.56	5.22	0.41	1.42	1.23	1.08	1.06	1.07	0.46	0.21	–.964
2003	2.79	7.99	14.70	33.22	35.01	5.84	0.44	1.25	1.34	1.04	1.09	1.07	0.46	0.30	–.857
2004	2.37	7.79	13.55	34.38	35.80	5.61	0.49	1.41	1.33	1.07	1.14	1.09	0.39	0.19	–.893
2005	2.49	7.55	13.73	34.55	35.12	5.87	0.70	1.29	1.31	1.19	1.16	1.07	0.38	0.24	–.964
2006	2.81	7.60	13.76	33.37	34.81	7.03	0.62	1.32	1.43	1.18	1.16	1.06	0.41	0.29	–.964
2007	3.43	7.25	15.89	30.83	34.67	7.30	0.62	1.48	1.25	1.37	1.15	1.07	0.39	0.27	–.964
2008	3.93	7.88	16.63	27.25	34.88	8.47	0.96	1.55	1.24	1.31	1.13	1.02	0.45	0.76	–.929
M	3.33	7.71	17.44	32.86	32.25	5.81	0.60	1.40	1.23	1.15	1.09	1.04	0.45	0.29	–1.000

cates a positive correlation between the workers’ age and the seriousness of the accident. This coincides with Bennet and Passmore’s results [11].

Table 5 shows the ADIs for the seven most frequent types of accidents in underground and surface energetic mining, according to the experience of the injured workers.

We calculated the Spearman rank correlation coefficient for each type of accident to evaluate the relationship between experience and the ADI. The null hypothesis can be rejected only for accident 50 (contact with a cutting, piercing, hard or rough material agent), for underground energetic mining. In the other kinds of accidents in underground and surface mining, the null hypothesis cannot be rejected because the Spearman rank

correlation coefficients are under the critical value of .786. So, for the other types of accidents, there is no statistically significant relationship between the injured parties and the seriousness of the accidents expressed through the ADI.

To compare the ADI for each group, considering experience in underground and surface energetic mining, we used the Mann–Whitney *U* test. The results showed the null hypothesis could not be rejected for accident 31 (blow, or hitting something as a result of a fall), 32 (blow as a result of a fall, or crashing into an immovable object), 71 (physical overexertion of the musculoskeletal system) and the remaining types of accidents (considered as one type). We concluded that there was no significant difference between

TABLE 3. Distribution of Lost Workdays and Nonfatal Accidents by Age Group (1999–2008)

Year	Lost Workdays							Total
	16–24	25–29	30–34	35–39	40–44	45–54	≥55	
1999	9635	18237	48436	85356	72940	16206	2313	253 123
2000	9544	18735	43832	73243	71067	17211	2078	235 710
2001	5033	14252	47890	89861	90407	15850	1544	264 837
2002	5178	13938	32095	69485	76711	12164	851	210 422
2003	4248	13455	23494	59635	66526	11623	863	179 844
2004	2270	9149	16241	46469	50023	8551	1087	133 790
2005	2779	8892	17048	35815	38206	7878	1031	111 649
2006	2411	5832	11563	27040	32702	6403	655	86 606
2007	1988	6247	15304	30394	30467	8785	1025	94 210
2008	2557	5458	12438	20520	25733	7828	1574	76 108
total	45643	114 195	268 341	537 818	554 782	112 499	13 021	1 646 299

Year	Nonfatal Accidents							Total
	16–24	25–29	30–34	35–39	40–44	45–54	≥55	
1999	495	955	2391	3859	3075	634	74	11 483
2000	464	847	2162	3329	3060	581	70	10 513
2001	235	614	1735	3101	2929	444	54	9 112
2002	215	557	1248	2462	2589	391	30	7 492
2003	182	520	959	2164	2283	381	29	6 518
2004	136	450	782	1983	2066	322	28	5 767
2005	114	345	628	1582	1606	268	32	4 575
2006	108	295	534	1295	1350	273	24	3 879
2007	133	280	615	1194	1344	281	24	3 871
2008	139	279	589	964	1234	298	34	3 537
total	2221	5142	11 643	21 933	21 536	3 873	399	66 747

TABLE 4. Distribution of Average Duration Index for Nonfatal Injuries by Age Group (1999–2008)

Year	Average Duration Index							Spearman Correlation
	16–24	25–29	30–34	35–39	40–44	45–54	≥55	
1999	19.46	19.10	20.26	22.12	23.72	25.56	31.26	.964
2000	20.57	22.12	20.27	22.00	23.22	29.62	29.69	.893
2001	21.42	23.21	27.60	28.98	30.87	35.70	28.59	.787
2002	24.08	25.02	25.72	28.22	29.63	31.11	28.37	.893
2003	23.34	25.88	24.50	27.56	29.14	30.51	29.76	.929
2004	16.69	20.33	20.77	23.43	24.21	26.56	38.82	1.000
2005	24.38	25.77	27.15	22.64	23.79	29.40	32.22	.464
2006	22.32	19.77	21.65	20.88	24.22	23.45	27.29	.714
2007	14.95	22.31	24.88	25.46	22.67	31.26	42.71	.893
2008	18.40	19.56	21.12	21.29	20.85	26.27	46.29	.893
total	20.55	22.21	23.05	24.52	25.76	29.05	32.63	1.000

TABLE 5. Average Duration Index for 7 Types of Accident-Related Injuries Depending on Experience in Underground and Surface Mining (2003–2008)

Experience in Underground Mining	Average Duration Index							All Types of Accidents
	71	42	50	32	40	41	31	
Nonfatal injuries	5084	4925	3874	1353	1317	882	840	23970
Fatalities	0	4	0	0	0	1	3	20
Lost workdays (nonfatal injuries)	120680	122033	80711	26647	37606	14498	22190	569774
Type of accident (total)	23.74	24.78	20.83	20.23	27.79	16.44	26.42	23.77
0–12 months	23.81	25.43	21.69	23.20	24.67	16.77	23.84	24.56
13–30 months	25.95	23.82	28.30	14.60	37.75	19.95	24.17	26.17
31–60 months	23.45	21.78	20.27	63.60	7.24	18.95	29.81	22.42
61–120 months	22.64	25.97	18.79	14.88	41.42	15.98	26.41	23.29
121–180 months	25.97	22.39	18.63	45.92	16.27	14.61	25.26	24.01
181–240 months	22.34	23.46	16.86	14.79	25.79	10.97	26.67	21.50
≥241 months	27.26	32.64	15.09	16.29	17.06	18.20	51.25	27.80
Spearman correlation	.143	.179	–.964	–.071	–.143	–.464	.750	–.071

Experience in Surface Mining	Average Duration Index							All Types of Accidents
	71	42	31	32	44	41	63	
Nonfatal Injuries	424	211	116	73	57	49	43	1372
Fatalities	0	1	0	0	0	0	0	3
Lost workdays (nonfatal injuries)	10266	6111	3264	1662	1286	660	1292	34513
Type of accident (total)	24.21	28.96	28.14	22.77	22.56	13.47	30.05	25.16
0–12 months	21.49	24.59	27.57	20.69	31.26	13.72	24.29	23.94
13–30 months	20.69	37.93	28.05	19.00	20.50	7.33	22.50	23.18
31–60 months	30.03	28.73	26.95	17.33	14.54	28.75	38.63	27.61
61–120 months	21.46	32.38	38.63	49.50	16.20	4.40	28.00	23.47
121–180 months	22.64	24.71	14.83	18.20	28.00	12.25	27.00	24.94
181–240 months	27.44	40.47	32.37	22.25	23.50	13.25	46.83	28.98
≥ 241 months	26.65	26.95	21.83	32.33	14.50	14.00	0.00	26.06
Spearman correlation	.464	.214	–.214	.393	–.429	.107	.071	.393

Notes. 31—blow, or hitting something as a result of a fall; 32—blow as a result of a fall, or crashing into an immovable object; 40—crashing into or hitting a moving object; 41—being hit by an object or projected fragments; 42—being hit by a falling object or one that is detached; 44—crash or blow against a moving object, including vehicles (immovable worker); 50—contact with a cutting, piercing, hard or rough material agent; 71—physical overexertion of the musculoskeletal system; 63—being trapped or flattened.

energetic underground and surface mining for those four types of accidents. This is not the case for accident 41 (being hit by an object or projected fragments) or 42 (being hit by a falling object or one that is detached), since the null hypothesis is rejected because in both cases the *U* values are under or equal to the critical value.

3.2. Size of the Mines

Table 6 shows the percentage of workers by year and size of mine, for 1999–2006. The workforce

is divided into six sizes of mines (in number of workers) classified in percentage per year.

Tables 6–7 show how the total number of accidents does not coincide exactly with the number of those in Table 2. This is so because in the classification according to size, the database field has missing values. Mining data for 2007–2008 were not published yet and, consequently, we could not calculate either the percentage of workers according to the size of the mines or the risk index for those years.

TABLE 6. Energetic Mining Population by Size of Mine (1999–2006)

Year	Size of Mine (%)					
	1–9	10–19	20–49	50–99	100–499	≥500
1999	0.24	0.79	3.77	4.31	21.64	69.25
2000	0.22	0.84	4.09	5.87	23.09	65.88
2001	0.20	1.20	3.94	4.73	21.25	68.68
2002	0.36	0.94	4.60	5.78	22.70	65.61
2003	0.32	1.18	3.89	6.20	22.32	66.10
2004	0.20	1.41	5.76	5.70	24.62	62.31
2005	0.31	1.78	3.35	6.83	33.63	54.09
2006	0.29	1.45	4.32	5.72	37.19	51.04
<i>M</i>	0.27	1.20	4.22	5.64	25.80	62.87

The results in Table 7 indicate a significantly greater risk for 1–9 workers (especially), 10–19, 20–49 and 50–99 workers, with the estimated risk indices of 3.33, 1.25, 1.26 and 1.32, respec-

tively. At the same time, risk was significantly lower for the 100–499 category, with an index of .76. It should be noted that 100–499 workers (especially) and ≥500 workers were the safest categories in 1999–2006. The 1–9 workers category was more dangerous because the risk index was significantly higher than for the other groups. These results coincide with studies that established that the proportion of accidents was greater in small mines [13, 20].

Spearman coefficients calculated for distributions of the risk index according to the size of the mines showed that there was no correlation between accidents and the size of the mines. The null hypothesis can be rejected for 1999 and 2002 only. For other years, Spearman rank correlation coefficients were high (except for 2004), but they

TABLE 7. Distribution of Accidents by Size of Mine and Estimated Risk Index (1999–2006)

Year	Size of Mine (%)						Risk Index						Spearman Correlation
	1–9	10–19	20–49	50–99	100–499	≥500	1–9	10–19	20–49	50–99	100–499	≥500	
1999	1.33	1.69	4.58	4.46	18.01	69.93	5.47	2.13	1.21	1.04	0.83	1.01	–.943
2000	1.33	1.80	4.16	7.46	17.80	67.45	6.13	2.13	1.02	1.27	0.77	1.02	–.786
2001	0.54	1.20	3.73	6.40	19.88	68.26	2.70	1.00	0.95	1.35	0.94	0.99	–.600
2002	0.75	1.59	5.74	6.18	18.19	67.55	2.05	1.68	1.25	1.07	0.80	1.03	–.943
2003	1.07	1.29	4.68	8.61	19.53	64.81	3.40	1.10	1.20	1.39	0.88	0.98	–.714
2004	0.71	1.04	5.06	8.96	18.62	65.62	3.58	0.74	0.88	1.57	0.76	1.05	–.143
2005	0.72	1.53	5.98	9.95	20.34	61.48	2.29	0.86	1.78	1.46	0.60	1.14	–.543
2006	0.67	1.86	8.74	7.71	24.74	56.29	2.34	1.28	2.02	1.35	0.67	1.10	–.771
<i>M</i>	0.89	1.50	5.33	7.47	19.64	65.17	3.33	1.25	1.26	1.32	0.76	1.04	–.714

TABLE 8. Distribution of Accidents and Lost Workdays by Size of Mine Categories (1999–2008)

Year	Lost Workdays						Nonfatal Accidents					
	1–9	10–19	20–49	50–99	100–499	≥500	1–9	10–19	20–49	50–99	100–499	≥500
1999	3348	4755	12308	10048	46903	141777	140	177	481	468	1885	7336
2000	3251	4732	9824	15472	43806	131631	129	175	403	722	1724	6547
2001	1510	3188	7823	13173	47536	179492	47	105	326	559	1738	5970
2002	1215	2475	9583	11269	37498	141554	54	115	416	448	1318	4894
2003	1757	1900	7992	12574	34458	118790	69	82	301	553	1255	4160
2004	1541	1415	7923	12592	29353	80966	41	60	291	516	1074	3785
2005	1341	2153	6543	11981	23769	65862	33	70	273	453	930	2816
2006	505	2200	7905	6577	24044	45348	26	72	338	299	959	2184
2007	577	3557	10343	8773	37819	33119	21	140	425	389	1344	1551
2008	745	2461	10014	8526	29533	24829	41	80	386	421	1178	1431
total	15790	28836	90258	110985	354719	963368	601	1076	3640	4828	13405	40674

were lower than the critical value (.829). All the coefficients were negative.

To analyse the possible relationship between the seriousness of the accidents and the size of the mines, we calculated the ADI for each group.

Spearman rank correlation coefficients were lower than the critical value, except in 2002 and 2005 (the two coefficients had different signs). This indicates that there is no correlation between the size of the mines (defined by the number of

TABLE 9. Distribution of Average Duration Index for Nonfatal Injuries by Size of Mine Categories (1999–2008)

Year	Average Duration Index						Spearman Correlation
	1–9	10–19	20–49	50–99	100–499	≥500	
1999	23.91	26.86	25.59	21.47	24.82	19.32	–.543
2000	25.20	27.04	24.32	21.34	25.32	20.08	–.543
2001	32.13	30.36	24.00	23.52	27.32	30.05	–.486
2002	22.50	21.52	23.04	25.15	28.45	28.92	.943
2003	25.46	22.89	26.55	22.74	27.46	28.53	.600
2004	37.59	23.58	27.13	24.36	27.31	21.37	–.486
2005	40.64	30.76	23.88	26.27	25.50	23.38	–.829
2006	19.42	30.56	23.32	22.00	25.05	20.76	.029
2007	27.48	25.41	24.34	22.55	28.14	21.35	–.429
2008	18.17	30.76	25.94	20.25	25.07	17.35	–.371
<i>M</i>	26.27	26.80	24.80	22.99	26.46	23.69	–.429

TABLE 10. Distribution of Accidents and Worked Hours by Size of Mine Category (1999–2006)

Year	Lost Workdays					
	1–9	10–19	20–49	50–99	100–499	≥500
1999	3348	4755	12308	10048	46903	141777
2000	3251	4732	9824	15472	43806	131631
2001	1510	3188	7823	13173	47536	179492
2002	1215	2475	9583	11269	37498	141554
2003	1757	1900	7992	12574	34458	118790
2004	1541	1415	7923	12592	29353	80966
2005	1341	2153	6543	11981	23769	65862
2006	505	2200	7905	6577	24044	45348
total	14468	22818	69901	93686	287367	905420

Year	Worked Hours					
	1–9	10–19	20–49	50–99	100–499	≥500
1999	79783	183880	839619	761356	4778611	12861750
2000	175422	164031	767758	1212769	4512380	11982640
2001	63654	204159	688551	769283	3740067	11577285
2002	76788	153575	715691	998240	3302615	9784091
2003	66338	152578	529969	841758	3125267	8831090
2004	39264	152694	674034	582418	3087323	7076267
2005	46661	298114	467478	966063	4711932	7262752
2006	46896	250110	591405	757276	5398199	6600115
total	594806	1559141	5274504	6889163	32656395	75975991

TABLE 11. Distribution of Gravity Index for Nonfatal Injuries by Size of Mine (1999–2006)

Year	Gravity Index						Spearman Correlation
	1–9	10–19	20–49	50–99	100–499	≥500	
1999	41.96	25.86	14.66	13.20	9.82	11.02	–.943
2000	18.53	28.85	12.80	12.76	9.71	10.99	–.886
2001	23.72	15.62	11.36	17.12	12.71	15.50	–.486
2002	15.82	16.12	13.39	11.29	11.35	14.47	–.543
2003	26.49	12.45	15.08	14.94	11.03	13.45	–.543
2004	39.25	9.27	11.75	21.62	9.51	11.44	–.314
2005	28.74	7.22	14.00	12.40	5.04	9.07	–.543
2006	10.77	8.80	13.37	8.69	4.45	6.87	–.771
<i>M</i>	24.32	14.63	13.25	13.60	8.80	11.92	–.886

workers) with the seriousness of the accidents. Table 9 shows the values of the ADI for six sizes of mines. Table 9 was calculated from the data in Table 8. There was no important variation between the different sizes of mines.

To obtain information about for the possible relationship between the seriousness of the accidents and the size of the mines, we calculated the gravity index for each group. Table 11 shows that index; it was developed from the data in Table 10.

Spearman correlation coefficients were lower than the critical value (.886), except in 1999, 2000, and the mean values in 1999–2006. This indicates a poor negative correlation between the size of the mine and the seriousness of the accidents, expressed as the gravity index. Furthermore, there is an important variation of the gravity index among the different sizes of mine. Thus, smaller mines had a greater proportion of accidents with more serious injuries. This coincides with Hunting and Weeks [13], Fabiano, Currò and Pastorino [19] and Saari [20].

4. CONCLUSIONS

We can draw the following conclusions.

- The incidence of work accidents in Spanish energetic mining decreases, whereas the age of the injured workers increases. However, the seriousness of the injuries caused by the accidents increases with age. Both results were significant. Using these results, the competent administrations and prevention services of the Spanish energetic mining sector should

programme specific safety training and information for the youngest workers (especially those under 29 years). The causes of the most serious accidents of the oldest workers should also be analysed (especially those over 54 years). A possible explanation could be that the older workers take longer to recover from the same injuries than the younger ones.

- We did not observe any relationship between the seriousness of the accidents and the experience of the injured parties. The analysis was carried out for the seven most frequent kinds of accidents in underground and surface mining. There was a relationship between the seriousness of the accidents and the workers' occupational experience in accident type 50 (contact with a cutting, piercing, hard or rough agent) in underground mining only. Thus, the negative consequences of the accidents produced by type 50 decrease, whereas the workers' occupational experience increases.
- Accident 71 (physical overexertion of the musculoskeletal system) and 42 (being hit by a falling object or one that is detached) are the most frequent in underground and surface mining. Those two types of accidents caused the highest number of lost workdays, specifically, 242713 and 16377 lost workdays in underground and surface mining, respectively. It should be taken into account that in underground mining there is the highest ADI by the group of workers with more experience (>240 months), with values of 32.64 and 27.26 for accidents 42 and 71, respectively. Thus, in energetic underground mining for 1999–2008,

for workers with over 240 months of occupational experience (usually 40–45 years old), the results showed that the injuries caused by overexertion produced more lost workdays by the oldest workers. This is why people who are responsible for the organization of the work in different mines (underground and surface) in the Spanish energetic mining sector should consider the age of the workers who have to carry out specific jobs involving physical effort.

- The results seem to show that mining centres with a low number of workers had a higher incidence of accidents than those with more workers. This incidence is especially important in mines with under 10 workers, with an average risk index of 3.33 in 1999–2006.
- We cannot confirm that the seriousness of the accidents, expressed as ADI, increases or decreases with the size of the mining work centres, due to low statistical significance. However, if the seriousness of the accidents is expressed as a gravity index, the results indicate that the accidents in small mines are more serious. Thus, for 1999–2006, the average gravity index for energetic mining was 24.32 and 11.92, which corresponds with mines with ≤ 9 and ≥ 500 workers, respectively.

REFERENCES

1. Government of Spain. Ministry of Employment and Social Security. Accidentes de trabajo con baja, según gravedad, por sector y rama de actividad [Accidents at work with lost workdays, by gravity, by sector and group of industry]. Retrieved March 21, 2012, from: http://www.meyss.es/estadisticas/eat/eat06/A1/a12_top_EXCEL.htm
2. Government of Spain. Ministry of Industry, Energy and Tourism. Estadística minera [Mining statistics]. Retrieved March 21, 2012, from: <http://www.minetur.gob.es/energia/mineria/Estadistica/Paginas/Consulta.aspx>
3. Government of Spain. Ministry of Employment and Social Security. Ocupados, según sexo y edad (1) [Employed workers by sex and age (1)]. Retrieved March 21, 2012, from: http://www.meyss.es/estadisticas/bel/EPA/epa8_top_EXCEL.htm
4. National Institute for Occupational Safety and Health (NIOSH). 1983–2007, number of mining employees, total number of hours worked, and number of full-time equivalent employees by year; excludes office employees. 2009. Retrieved March 21, 2012, from: <http://www.cdc.gov/niosh/mining/statistics/tables/e2.html>
5. Queensland Mines and Quarries. Safety Performance and Health Report. City East, QLD, Australia: Queensland Government; 2011. Retrieved March 21, 2012, from: http://mines.industry.qld.gov.au/assets/health-report/health_report_2009-10.pdf
6. Murphy JN. Coal mine health and safety research in the USA—the achievements of the US Bureau of Mines. *Coal International*. 1994;242(6):219–26.
7. McKenna FP. Accident proneness: a conceptual analysis. *Accid Anal Prev*. 1983;15(1):65–71.
8. Sanmiquel L, Freijo M, Edo J, Rossell JM. Analysis of work related accidents in the Spanish mining sector from 1982–2006. *J Safety Res*. 2010; 41(1):1–7.
9. National Research Council. Toward safe underground coal mines. Washington, DC, USA: National Academy Press; 1982. Retrieved March 21, 2012, from: http://www.archive.org/stream/towardsaferunder024710mbp/towardsaferunder024710mbp_djvu.txt
10. Bennett JD. Relationship between workplace and worker characteristics and severity of injuries in U.S. underground bituminous coalmines, 1975–1981 [doctoral dissertation]. University Park, USA: Pennsylvania State University; 1982.
11. Bennett JD, Passmore DL. Multinomial logit analysis of injury severity in U.S. underground bituminous coal mines. *Accid Anal Prev*. 1985;17(5):399–408.
12. Butani SJ. Relative risk analysis of injuries in coal mining by age and experience at present company. *J Occup Accid*. 1988;10(3):209–16.

13. Hunting KL, Weeks JL. Transport injuries in small coalmines: an exploratory analysis. *Am J Ind Med.* 1993;23:391–406.
14. Phiri J. Development of statistical indices for the evaluation of hazards in Longwall Face Operations [doctoral dissertation]. University Park, USA: Pennsylvania State University; 1989.
15. Hansen CP. A causal model of the relationship among accidents, biodata, personality and cognitive factors. *J Appl Psychol.* 1989;74:81–90.
16. Leigh J, Mulder HB, Want GV, Farnsworth NP, Morgan GG. Personal and environmental factors in coal mining accidents. *J Occup Accid.* 1990;13(3): 233–50.
17. Maiti J, Bhattacharjee A. Evaluation of risk of occupational injuries among underground coalmine workers through multinomial logit analysis. *J Safety Res.* 1999;30(2):93–101.
18. Groves WA, Kecojevic VJ, Komljenovic D. Analysis of fatalities and injuries involving mining equipment. *J Safety Res.* 2007;38(7):461–70.
19. Fabiano B, Currñ F, Pastorino R. A study of the relationship between occupational injuries and firm size and type in the Italian industry. *Saf Sci.* 2004;42(7):587–600.
20. Saari J. Pequeñas y medianas empresas [Small and medium enterprises]. In: *Jornada técnica: La prevención de riesgos en la PYME.* Asepeyo; 2005. p. 11–20. Retrieved March 21, 2012, from: [http://www.asepeyo.es/web%5CBiblioteca.nsf/ficheros/monografia28abril2005.pdf/\\$file/monografia28abril2005.pdf](http://www.asepeyo.es/web%5CBiblioteca.nsf/ficheros/monografia28abril2005.pdf/$file/monografia28abril2005.pdf)